

**GAMBARAN KUNJUNGAN ANC PADA IBU HAMIL KEKURANGAN
ENERGI KRONIS DI PUSKESMAS MINGGIR
SLEMAN YOGYAKARTA
2012**

INTISARI

Agnes Indah Pramita¹, Ratih Kumoro Jati, S.SiT.,M.Kes², Nur Rahmawati Sholihah, SST³

Latar Belakang : Prevalensi KEK tingkat nasional untuk wanita berusia 15-45 tahun masih mencapai angka 13,6% dengan persentasi perkotaan dan pedesaan masing-masing adalah 13,0% dan 14,1% (Depkes 2007). Kunjungan Antenatal Care (ANC) adalah kunjungan ibu hamil ke bidan atau dokter sedini mungkin semenjak ia merasa dirinya hamil untuk mendapatkan pelayanan/asuhan antenatal. KEK merupakan keadaan satus gizi kurang pada ibu hamil. Penyebab KEK belum diketahui secara pasti, namun penyebab utama dikarenakan karena kurangnya asupan energi dan protein. Menurut Wiknjosatiro (2008), faktor-faktor yang mempengaruhi terjadinya KEK diantaranya adalah kepatuhan kunjungan ANC, pendapatan keluarga dan asupan makanan.

Tujuan Penelitian : Mengetahui gambaran kunjungan ANC pada ibu hamil yang KEK di Puskesmas Minggir, Sleman, Yogyakarta 2012.

Metode Penelitian : Penelitian ini menggunakan metode *deskriptif* dengan pendekatan *crosssectional*. Tehnik sampel yang digunakan adalah *total sampling* dengan jumlah sampel 58 ibu hamil KEK. Tehnik analisis yang dipakai dalam penelitian ini dengan analisis data deskriptif.

Hasil Penelitian : Frekuensi kunjungan ANC dikategorikan baik sebanyak 25 orang (43,1%) dan dikategorikan kurang sebanyak 33 orang (56,9%). Sebagian besar responden ibu termasuk usia 20 – 35 tahun, yaitu ada 53 responden (91,4%), pendidikan sampai tingkat SMA, yaitu ada 32 responden (55,2%), paritas primipara, yaitu ada 26 responden (44,8%) dan ibu yang tidak bekerja (IRT), yaitu ada 27 responden (46,6%).

Kesimpulan : Frekuensi kunjungan ANC dikategorikan baik sebanyak 25 orang (43,1%) dan dikategorikan kurang sebanyak 33 orang (56,9%).

Kata Kunci : Gambaran Kunjungan ANC Pada Ibu Hamil Kekurangan Energi Kronis

Halaman : i – xiv, 1-43

Pustaka : 44 buku (2002-2010), 4 KTI

¹Mahasiswa Diploma III Kebidanan STIKES A. Yani Yogyakarta

²Dosen STIKES A. Yani Yogyakarta

³Dosen STIKES A. Yani Yogyakarta

**DESCRIPTION VISIT TO PREGNANT WOMEN ANC ENERGY
SHORTAGE IN CHRONIC HEALTH MOVE OVER SLEMAN
YOGYAKARTA
2012**

ABSTRACT

Agnes Indah Pramita¹, Ratih Kumoro Jati, S.SiT.,M.Kes², Nur Rahmawati Sholihah, SST³

Background: KEK national prevalence rate for women aged 15-45 years still reached 13.6% as a percentage of urban and rural areas respectively 13.0% and 14.1% (Ministry of Health 2007). According Sadli and Bakhtiar (2010), KEK is one of the causes of high maternal mortality rate in Indonesia in addition to hemorrhage, eclampsia, abortion, obstructed labor and infection. Visits Antenatal Care (ANC) is a pregnant mother's visit to the midwife or doctor as early as possible since she felt herself to be pregnant to get care / antenatal care. KEK is the status of the state of malnutrition among pregnant women. Cause of SEZ yet known with certainty, but the main cause is because due to lack of energy and protein intake. According Wiknjosatno (2008), the factors that influence the occurrence of which is compliance KEK ANC visits, family income and food intake.

Objective: Knowing overview ANC visits in pregnant women in health centers KEK Move, Sleman, Yogyakarta 2012

Methods: This study uses a descriptive cross sectional approach. Sampling technique used is total sampling with a sample of 58 pregnant women KEK. Tehnik analysis used in this study with descriptive data analysis.

Result : The results of this study showed that the frequency of ANC visits were categorized either as many as 25 people (43.1%) and categorized approximately 33 persons (56.9%)

Conclusion: The majority of respondents, including maternal age 20-35 years, ie there were 53 respondents (91.4%), education up to high school level, there are 32 respondents (55.2%), primiparous parity, ie there are 26 respondents (44, 8%) and mothers who do not work (IRT), ie there were 27 respondents (46.6%). Most of the respondents have less frequency of ANC visits category, ie there are 33 respondents with the percentage of 56.9%.

Keywords: An ANC Visits In Chronic Energy Deficiency Pregnant Women

Pages : i-xiv, 1-43

Library : 44 book(2002-2010),4 KTI

¹ Midwifery Diploma Students STIKES A. Yani Yogyakarta

² Lecturer STIKES A. Yani Yogyakarta

³ Lecturer STIKES A. Yani Yogyakarta