

**GAMBARAN IBU HAMIL DENGAN KEJADIAN PENYULIT
PERSALINAN DI PUSKESMAS GAMPING 1**

SLEMAN YOGYAKARTA TAHUN 2013

Niken Damayanti¹, Ekawati²

¹Mahasiswa DIII Prodi Kebidanan STIKes Jenderal Achmad Yani Yogyakarta

²Dosen STIKes Jenderal Achmad Yani Yogyakarta

INTISARI

Latar Belakang: Penyebab kematian maternal dari faktor reproduksi diantaranya adalah *maternal age*/usia ibu. AKI di Indonesia meningkat menjadi 359/100.000 kelahiran hidup. Komplikasi lain mungkin ada sebelum kehamilan tetapi memburuk selama kehamilan. Komplikasi utama yang menjelaskan 80% dari seluruh kematian ibu adalah: perdarahan parah (terutama pendarahan setelah melahirkan), Infeksi (biasanya setelah melahirkan), tekanan darah tinggi selama kehamilan (pre-eklampsia dan eklampsia), tidak aman aborsi

Tujuan Penelitian: Untuk mengetahui gambaran ibu hamil dengan kejadian penyulit persalinan di Puskesmas Gamping 1 Sleman Yogyakarta Periode Januari-Desember 2013.

Metode Penelitian: Penelitian ini merupakan jenis penelitian *diskriptif non analitik*. Lokasi penelitian di Puskesmas Gamping 1 Sleman Yogyakarta. Sampel dalam penelitian sebanyak 77 responden yang mengalami penyulit persalinan dengan teknik sampling jenuh. Alat pengumpulan data adalah data rekam medik. Teknik analisa data adalah analisa univariat dengan program SPSS.

Hasil: Jumlah ibu hamil dengan penyulit persalinan sesuai kelompok usia yaitu usia <20 tahun sebanyak 18 ibu (23.4%), usia reproduksi sehat 20-35 tahun sebanyak 14 ibu (18.2%), usia >35 tahun sebanyak 45 (58.4%). Jumlah ibu hamil dengan paritas tertinggi yaitu pada multipara sebanyak 47 ibu (61.0) dan terendah pada primipara sebanyak 30 ibu (39.0). Jumlah ibu hamil dengan jenis persalinan seperti persalinan secara Spontan sejumlah 68 (88.3%), secara sectio caesarea sejumlah 9 (11.7%). Jumlah ibu hamil dengan jenis penyulit persalinan terbanyak pada hiperetnsi sejumlah 35 ibu (45.5%) dan terendah pada posisi saat bersalin sebanyak 2 ibu (2.6%).

Kesimpulan: Gambaran ibu hamil dengan kejadian penyulit persalinan di Puskesmas Gamping 1 Sleman Yogyakarta dapat di tarik kesimpulan bahwa ibu hamil yang mengalami penyulit persalinan ad 77 responden.

Kata Kunci: Ibu hamil dengan kejadian penyulit persalinan.

**DESCRIPTION OF PREGNANT MATERNAL WITH POSTPARTUM
COMPLICATIONS INCIDENTS IN PUSKESMAS GAMPING 1
SLEMAN YOGYAKARTA YEAR 2013**

Niken Damayanti¹, Ekawati²

¹High student of D-III Midwife STIKES Jenderal Achmad Yani Yogyakarta

²Lecturer of STIKES Jenderal Achmad Yani Yogyakarta

ABSTRACT

Background: The causes of maternal deaths from reproductive factors include maternal age / age of the mother. AKI in Indonesia increased to 359 / 100,000 live births. Other complications may exist before pregnancy but deteriorated during pregnancy. The main complication that explained 80% of all maternal deaths are: severe bleeding (mostly bleeding postpartum), infections (usually after giving birth), high blood pressure during pregnancy (pre-eclampsia and eclampsia), unsafe abortion.

Objective: To determine the description of pregnant maternal with postpartum complications incidents at Puskesmas Gamping 1 Sleman Yogyakarta Period of January to December, 2013.

Methods: This study is a descriptive study non-analytic types. The research sites in Puskesmas Gamping 1 Sleman Yogyakarta. The sample in the study a total of 77 respondents who experienced complications postpartum with the saturation sampling technique. Data collection tool was a medical record. Data analysis technique is a univariate analysis using SPSS software.

Results: The number of pregnant maternal with complications postpartum which appropriate age group are age < 20 years as many as 18 maternal (23.4%), healthy reproductive age 20 - 35 years as many as 14 maternal (18.2%), age > 35 years were 45 maternal (58.4%). The number of pregnant maternal with the highest parity on multiparous as many as 47 maternal (61.0%) and lowest in primiparous as many as 30 maternal (39.0%). The number of pregnant maternal with the type of postpartum like for spontaneous postpartum as many as 68 maternal (88.3%), as sectio caesarea as many as 9 maternal (11.7%). The number of pregnant maternal with the type of postpartum complications most in hypertension as many as 35 maternal (45.5%) and lowest in the position of during postpartum as many as 2 maternal (2.6%).

Conclusion: Description of pregnant maternal with postpartum complications incidents in Puskesmas Gamping 1 Sleman Yogyakarta can be deduced conclusion that pregnant maternal who experience postpartum complications as many as 77 respondents.

Keywords: Pregnant maternal with postpartum complications incidents.