

PERBEDAAN ONSET PENGELUARAN KOLOSTRUM PADA IBU POST PARTUM DENGAN PERSALINAN NORMAL DAN *SECTIO CAESARIA* DI RSUD PANEMBAHAN SENOPATI BANTUL

Maria NSW Kause¹ , Yanita Trisetiyaningsih² , Anastasia Suci Sukmawati³

INTISARI

Latar Belakang : Ibu yang mengalami operasi *sectio caesaria* akan merasa sakit, nyeri, dan efek penggunaan anastesi yang berkepanjangan sehingga belum bisa untuk melakukan Insiasi Menyusu Dini dibandingkan ibu dengan persalinan normal serta penggunaan anastesi yang memperlambat onset laktasi. Ibu yang melakukan persalinan normal akan mengalami onset laktasi yang cepat dan lebih mudah dikarenakan tidak mengalami nyeri dan langsung melakukan Inisiasi Menyusu Dini,

Tujuan Penelitian : Penelitian ini bertujuan untuk mengetahui perbedaan onset pengeluaran kolostrum pada ibu post partum dengan persalinan normal dan *sectio caesaria*.

Metode Penelitian : Rancangan penelitian ini adalah *cross sectional*, dengan jumlah sampel sebanyak 80 responden dengan teknik *purposive sampling*. Instrumen penelitian dalam penelitian ini adalah lembar observasi. Analisis data menggunakan uji *Mann-Whitney*.

Hasil : Dengan rata-rata waktu pada ibu post partum dengan persalinan normal adalah 13,6 jam dan pada ibu post partum dengan persalinan *sectio caesaria* adalah 22,6 jam. Hasil uji *Mann-Whitney* yaitu terdapat perbedaan onset pengeluaran kolostrum pada ibu post partum dengan persalinan normal dan *sectio caesaria* dengan nilai $p = 0.001 (p < 0,05)$ artinya H_0 ditolak.

Kesimpulan : Terdapat perbedaan onset pengeluaran kolostrum pada ibu post partum dengan persalinan normal dan *sectio caesaria*.

Kata Kunci : kolostrum, Persalinan normal , Persalinan *sectio caesaria*.

¹Mahasiswa S1 Keperawatan Stikes Jenderal Achmad Yani Yogyakarta

²Dosen PSIK Stikes Jenderal Achmad Yani Yogyakarta

³Dosen PSIK Stikes Jenderal Achmad Yani Yogyakarta

THE DIFFERENCE BETWEEN COLOSTRUM RELEASING ONSET IN POST PARTUM MOTHERS WITH NORMAL LABORS AND SECTIO CAESARIA IN RSUD PANEMBAHAN SENOPATI BANTUL

Maria NSW Kause¹ , Yanita Trisetiyaningsih² , Anastasia Suci Sukmawati³

ABSTRACT

Background : Mothers who undergo sectio caesarea surgery will experience illness, pain, and effects due to prolonged usage of anesthesia so that mothers cannot initiate early breastfeeding compared to mothers who undergo normal labor and usage of anesthesia which delays onset of lactation. Mothers who undergo normal labor will experience earlier and easier onset of lactation due to the absence of pain and immediate initiation of early breastfeeding.

Objective : To find out the difference between colostrum releasing onset in post partum mothers with normal labors and sectio caesaria in Panembahan Senopati General Hospital, Bantul.

Method : The design of this study was cross sectional. The number of samples were 80 respondents selected with purposive sampling technique. Study instrument was observational sheets. Data analysis applied Mann-Whitney test.

Result : The average time duration for post partum mothers with normal labor was 13,6 hours and for post partum mothers with sectio caesaria surgery was 22,6 hours. The result of Mann-Whitney test revealed that there was a difference between the onset of colostrum outing release in post partum mothers with normal labor and post partum mothers with sectio caesaria surgery with p value 0,001 ($p < 0,05$) which meant that H_0 was rejected.

Conclusion : There was a difference of colostrum outing release between post partum mothers with normal labor and with sectio caesaria surgery.

Keyword : Colostrum, Normal labors, Sectio Caesaria labors.

¹Student of Nursing Science Program on Collage of Health science General Achmad Yani Yogyakarta

²Lecturer of Nursing Science Program on Collage of Health science General Achmad Yani Yogyakarta

³Lecturer of Nursing Science Program on Collage of Health science General Achmad Yani Yogyakarta