

TINGKAT PENGETAHUAN IBU HAMIL TENTANG TANDA BAHAYA ANEMIA PADA IBU HAMIL TRIMESTER I DAN TRIMESTER III DI PUSKESMAS KASIHAN I BANTUL YOGYAKARTA

Risa Novi Yana Yanti¹, Ika Fitria Ayuningtyas²

INTISARI

Latar Belakang: Anemia merupakan salah satu faktor resiko terjadinya kematian ibu melahirkan. Anemia berpengaruh kurang baik bagi ibu, baik dalam kehamilan, persalinan, nifas dan masa selanjutnya. Berbagai penyulit persalinan ditimbulkan dari kejadian anemia dalam kehamilan. Salah satu faktor yang diduga sebagai penyebab utama tingginya angka kejadian anemia ibu hamil Puskesmas Kasihan 1 Bantul adalah kurangnya pengetahuan ibu hamil tentang tanda bahaya anemia, sehingga ibu tidak menyadari bahwa dirinya mengalami anemia selama masa kehamilan dan tidak segera melakukan upaya pencegahan serta pengobatan.

Tujuan: Diketuinya tingkat pengetahuan Ibu hamil tentang tanda bahaya anemia pada ibu hamil Trimester I dan III di Puskesmas Kasihan 1 Bantul tahun 2013.

Metode Penelitian: Rancangan penelitian ini adalah survey dengan *incidental sampling*. Populasi 34 orang dan jumlah sampel 34 orang, analisis data uji deskriptif frekuentif.

Hasil Penelitian: Mayoritas pengetahuan ibu hamil TM I dan TM III tentang penyebab anemia adalah cukup sebanyak 21 orang (61,8%), tentang tanda dan gejala anemia mayoritas cukup 26 orang (76,5%), tentang cara pencegahan dan penanggulangan anemia mayoritas cukup 26 orang (76,5%).

Kesimpulan: Mayoritas tingkat pengetahuan ibu hamil tentang tanda bahaya anemia pada ibu hamil TM I dan TM III di Puskesmas Kasihan I Bantul Yogyakarta dalam kategori cukup 28 orang (82,4%).

Kata kunci: Tingkat pengetahuan tentang tanda bahaya anemia, ibu hamil TM I dan TM III

¹ Mahasiswa D-III Kebidanan STIKES A. Yani Yogyakarta

² Dosen Pembimbing DIII Kebidanan STIKES A. Yani Yogyakarta

**KNOWLEDGE LEVEL OF PREGNANT WOMEN ABOUT ANEMIA
DANGER SIGN AT TRIMESTER I AND TRIMESTER III PREGNANT
WOMEN IN PITY THE HEALTH CENTER AT
1 BANTUL YOGYAKARTA**

Risa Novi Yana Yanti¹, Ika Fitria Ayuningtyas²

ABSTRACT

Background: Anemia is a risk factor for maternal mortality. Anemia adversely affects the mother, both in pregnancy, childbirth, postpartum and subsequent period. Various delivery complications arising from anemia in pregnancy. One factor that is thought to be the main cause of the high incidence of maternal anemia Pity 1 Bantul Health Center is a lack of knowledge of pregnant women about the danger signs of anemia, so she does not realize that he had anemia during pregnancy and did not immediately take steps to prevent as well as cure.

Objective: Knowing the level of knowledge of pregnant women about the danger signs of anemia in pregnant women in the third trimester and I pity 1 Bantul Health Center in 2013.

Methods: The study design was a survey by incidental sampling. Population of 34 people and the number of samples of 34 people, frekuentif descriptive analysis of the test data.

Results: The majority of maternal knowledge of TM I and TM III of the causes of anemia are quite as many as 21 people (61.8%), about the signs and symptoms of anemia sufficient majority 26 (76.5%), about how to prevent and control anemia majority pretty 26 people (76.5%).

Conclusion: The majority of the level of knowledge of pregnant women about danger signs of anemia in pregnant women TM I and TM III in Bantul Yogyakarta Puskesmas I Pity the category of pretty 28 people (82.4%).

Keywords: Level of knowledge about danger signs of anemia, maternal TM I and TM III

¹Midwifery student STIKES D-III A. Yani Yogyakarta.

²Midwifery Lecturer DIII STIKES A. Yani Yogyakarta.