

ABSTRACT

Sri Windi Lestari¹, Endang Suprapti²

Background : One of the biggest causes of maternal death is postpartum hemorrhage, mainly in the third stage of labor. Postpartum hemorrhage can be prevented by handling the active management of the third stage of the appropriate standard of care of normal deliveries. Results of preliminary studies in hospitals Panembahan Senopati Bantul Yogyakarta from 15 midwives by 2 midwives (13.33%) indicated that lack of time management in oxytocin injection.

Objective : This study aims to determine the Active Management Implementation Overview Kala III In Hospital Panembahan Senopati Bantul Yogyakarta in 2013.

Methods of Research : The research method uses descriptive exploratory study using One-Shot approach. The samples in this study is the total sampling. Subjects were all midwives in the hospital delivery room Panembahan Senopati Bantul which totaled 15 people. Univariant analysis of the data presented by frequency distribution.

The Results : The results showed that the majority of midwives (73.33%) in Panembahan Senopati Bantul Hospital has implemented active management of the third stage in honor of normal delivery care standards.

Conclusion : Overview of the third stage of implementation of active management by midwives in hospitals Panembahan Senopati Bantul Yogyakarta Year 2013 has been carried out in accordance with good standards of normal delivery care (APN).

Keywords : Midwives, Active management of the third stage.

¹ Student Diploma III Midwifery STIKES A.Yani Yogyakarta

² Supervisor STIKES A.Yani Yogyakarta

INTISARI

Sri Windi Lestari³, Endang Suprapti⁴

Latar Belakang : Salah satu faktor terbesar penyebab kematian ibu adalah perdarahan pasca persalinan, terutama terjadi pada kala III persalinan. Perdarahan pasca persalinan dapat dicegah dengan penanganan manajemen aktif kala III yang sesuai standar asuhan persalinan normal. Hasil studi pendahuluan di RSUD Panembahan Senopati Bantul Yogyakarta yang dari 15 bidan sebanyak 2 bidan (13,33%) menunjukkan bahwa kurangnya manajemen waktu dalam penyuntikan oksitosin.

Tujuan : Penelitian ini bertujuan untuk mengetahui Gambaran Pelaksanaan Manajemen Aktif Kala III Di RSUD Panembahan Senopati Bantul Yogyakarta Tahun 2013.

Metode Penelitian : Metode penelitian menggunakan penelitian deskriptif yang bersifat eksploratif dengan menggunakan pendekatan *One-Shot*. Pengambilan sampel dalam penelitian ini adalah *total sampling*. Subyek penelitian adalah seluruh bidan di ruang bersalin RSUD Panembahan Senopati Bantul yang berjumlah 15 orang. Analisis data secara univariant yang disajikan dengan distribusi frekuensi.

Hasil : Hasil penelitian menunjukkan bahwa mayoritas bidan (73,33%) di RSUD Panembahan Senopati Bantul telah melaksanakan manajemen aktif kala III sesuai dengan standar asuhan persalinan normal.

Kesimpulan : Gambaran Pelaksanaan manajemen aktif kala III oleh bidan di RSUD Panembahan Senopati Bantul Yogyakarta Tahun 2013 sudah dilaksanakan dengan baik sesuai standar asuhan persalinan normal (APN).

Kata kunci : Bidan, Manajemen aktif kala III

³ Mahasiswa Diploma III Kebidanan STIKES A.Yani Yogyakarta

⁴ Dosen Pembimbing STIKES A.Yani Yogyakarta